



Canadian Poultry Consultants Ltd.

### **Broiler Farm Enhanced C&D**

1. Remove manure, litter and other loose debris from the house to a site as far away from the barn as possible. These materials contain disease carrying insects, bacteria or viruses that were shed by the previous flock.
2. Apply fly, beetle and rodent control at the farm to reduce their movement and breeding activity.
3. Remove all the residual feed as it is an attractive food source for the rodents and beetles, which can act as vectors for various pathogens.
4. Sweep/blow down the house from top to bottom. Remember to clean the fan blades, heaters, louvers, lighting fixtures, walls, other equipment, etc. Remember to clean mortality buckets, service areas/anterooms. Ensure all loose, organic material has been removed.
5. Clean the water lines with a suitable product and subsequently flush and drain them.
6. Wash with a high pressure washer, hot water, and detergent (ceiling, walls, feeders, drinkers, and other equipment). Cover all surfaces with the cleaning solution and allow the surfaces to soak. Hand-wash any sensitive electrical devices and switches.
7. Disinfect all cleaned and dried surfaces with a disinfectant, using the manufacturers' recommendations. Thermo-fogging is recommended.

### **Product Examples:**

- **Detergent:** *BIOGEL*
- **Disinfectant:** *Virocid, Profilm*
- **Water line Sanitation:** *Proxy-clean, CID 2000*
- **Insecticide:** *Tempo, Malathion*
- **Rodenticide:**
  - **Inside:** Bromethalin (*Rampage*)
  - **Outside:** Anticoagulants (*Jaguar, Hawk, Hombre*)